

CP violation in $B_s \rightarrow J/\psi\phi$ in ATLAS


Radek Novotný
on behalf of the **ATLAS** collaboration

Beauty 2019, Ljubljana
September 30, 2019





- ▶ $B_s^0 \rightarrow J/\psi \phi$ is used to measure CP-violation phase Φ_s potentially sensitive to New Physics
- ▶ In SM ϕ_s is related to the CKM elements and predicted with high precision

$$\Phi_s \simeq 2 \arg[-(V_{ts} V_{tb}^*)/(V_{cs} V_{cb}^*)] = -0.0363_{-0.0015}^{+0.0016} \text{ rad}$$




- ▶ Other quantity in B_s^0 mixing is $\Delta\Gamma_s = \Gamma_s^L - \Gamma_s^H$, where Γ_s^L and Γ_s^H are the decay widths of the different mass eigenstates. $\Delta\Gamma_s$ is not sensitive to New Physics, however measurement is interesting to test a theory.
- ▶ The New Physics processes could introduce additional contributions to the box diagrams describing the B_s^0 mixing



- Inner Detector: PIX, SCT and TRT, $p_T > 0.4 \text{ GeV}$, $|\eta| < 2.5$
 - Run2: new IBL 25% improvement of time resolution with respect to Run1.
 - time resolution remains stable within increasing pileup in Run 2
- Muon Spectrometer: triggering ($|\eta| < 2.4$), precision tracking ($|\eta| < 2.7$)

- Events collected with mixture of triggers based on $J/\psi \rightarrow \mu^+ \mu^-$ identification, with muon p_T thresholds of either 4 GeV or 6 GeV (vary over run periods)
- No lifetime or impact parameter cut at HLT level



Data:

- ▶ 80.5 fb⁻¹ of 13 TeV pp collision data from 2015-2017
- ▶ Statistically combined with Run1 ATLAS results:
 - ▶ 4.9 fb⁻¹ (7 TeV, pp 2011)
 - ▶ 14.3 fb⁻¹ (8 TeV, pp 2012)

MC samples:

- ▶ Signal $B_s^0 \rightarrow J/\psi \phi$ MC events
- ▶ MC samples for peaking backgrounds $B_d^0 \rightarrow J/\psi K^{*0}$, $B_d^0 \rightarrow J/\psi K \pi$, $\Lambda_b^0 \rightarrow J/\psi K p$
- ▶ MC samples for tagging calibration channel $B^\pm \rightarrow J/\psi K^\pm$
(systematics and cross-checks only, real data used for calibration)

Event:

- ▶ Triggers (previous slide) and good quality data
- ▶ At least one PV formed from at least 4 ID tracks
- ▶ At least one pair of ID+MS identified $\mu^+\mu^-$

 $J/\psi \rightarrow \mu^+\mu^-$

- ▶ Dimuon vertex fit $\chi^2/d.o.f. < 10$
- ▶ Three dimuon invariant mass windows for BB/BE/EE (barrel, endcap) muon combinations

 $\phi \rightarrow K^+K^-$

- ▶ $p_T(K) > 1 \text{ GeV}$
- ▶ $1008.5 \text{ MeV} < m(KK) < 1030.5 \text{ MeV}$

 $B_s^0 \rightarrow J/\psi(\mu^+\mu^-)\phi(K^+K^-)$

- ▶ $p_T(B_s^0) > 10 \text{ GeV}$
- ▶ Four-track vertex fit $\chi^2/d.o.f. < 3$ (J/ψ mass constrained)
- ▶ Keep only the candidate with best vertex fit $\chi^2/d.o.f.$ in event
- ▶ $5150 \text{ MeV} < m(B_s^0) < 5650 \text{ MeV} \rightarrow$ in total 3 210 429 B_s^0 candidates

- ▶ $B_s^0 \rightarrow J/\psi \phi$ = pseudoscalar to vector-vector
- ▶ Final state: admixture of CP -odd ($L = 1$) and CP -even ($L = 0, 2$) states
- ▶ Distinguishable through time-dependent angular analysis
- ▶ Non-resonant S -wave decay $B_s^0 \rightarrow J/\psi K^+ K^-$ contribute to the final state
- ▶ Included in the differential decay rate due to interference with the $B_s^0 \rightarrow J/\psi(\mu^+ \mu^-)\psi(K^+ K^-)$ decay




Figure: Angles between final state particles in transversity basis.

We perform unbinned maximum likelihood fit simultaneously for B_s^0 mass, decay time and the decay angles:

$$\begin{aligned} \ln \mathcal{L} = & \sum_{i=1}^N \{ w_i \cdot \ln(f_s \cdot \mathcal{F}_s(m_i, t_i, \sigma_m, \sigma_t, \Omega_i, P(B|Q), p_{\Gamma_i}) \\ & + f_s \cdot f_{B_d^0} \cdot \mathcal{F}_{B_d^0}(m_i, t_i, \sigma_m, \sigma_t, \Omega_i, P(B|Q), p_{\Gamma_i}) \\ & + f_s \cdot f_{\Lambda_b} \cdot \mathcal{F}_{\Lambda_b}(m_i, t_i, \sigma_m, \sigma_t, \Omega_i, P(B|Q), p_{\Gamma_i}) \\ & + (1 - f_s \cdot (1 + f_{B_d^0} + f_{\Lambda_b})) \cdot \mathcal{F}_{\text{bkg}}(m_i, t_i, \sigma_m, \sigma_t, \Omega_i, P(B|Q), p_{\Gamma_i})) \} \end{aligned}$$

Physics parameters

- ▶ CPV phase ϕ_s
- ▶ Decay widths: $\Delta\Gamma_s, \Gamma_s$
- ▶ Decay amplitudes: $|A_0(0)|^2, |A_{\parallel}(0)|^2, \delta_{\parallel}, \delta_{\perp}$
- ▶ S-wave: $|A_S(0)|^2, \delta_S$
- ▶ δm_s fixed to PDG

Observables

- ▶ Base observables : m_i, t_i, Ω_i
- ▶ Conditional observables per-candidate:
 - ▶ resolutions: $\sigma_{m_i}, \sigma_{t_i}$ ($B - p_{\Gamma_i}$ dependent)
 - ▶ tagging probability and method: $P(B|Q)$

$$\ln \mathcal{L} = \sum_{i=1}^N \left\{ \overset{\text{Tau weight}}{\underset{|}{w_i}} \cdot \ln \left(\overset{\text{Signal}}{\underset{|}{f_s \cdot \mathcal{F}_s}} + \overset{\text{Peaking background}}{\underset{|}{f_s \cdot f_{B_d^0} \cdot \mathcal{F}_{B_d^0} + f_s \cdot f_{\Lambda_b} \cdot \mathcal{F}_{\Lambda_b}}} + \overset{\text{Combinatorial background}}{\underset{|}{(1 - f_s \cdot (1 + f_{B_d^0} + f_{\Lambda_b})) \cdot \mathcal{F}_{\text{bkg}}}} \right) \right\}$$

- ▶ Data are corrected by the decay time correction
- ▶ Mass as well as lifetime use per-candidate width and scale factor, with flavour-dependent terms weighted by tagging probability $P(B|Q)$
- ▶ Contributions from $B_d^0 \rightarrow J/\psi K^{*0}$, $B_d^0 \rightarrow J/\psi K\pi$ and $\Lambda_b^0 \rightarrow J/\psi Kp$ due to wrong mass assignment (KK)
 - ▶ Efficiencies and acceptance from MC
 - ▶ BR from PDG
 - ▶ Fragmentation fractions from other measurements
- ▶ Combinatorial background for angular distribution use Legendre polynomials from sidebands; fixed in the main fit

▶ Opposite side tagging

- ▶ Use $b - \bar{b}$ pair correlation to infer initial signal flavour from the other B meson
- ▶ Provide the probability of signal candidate to be B_s^0 or \bar{B}_s^0

$$\begin{aligned} \mathcal{F}_s(m_i, t_i, \sigma_{t_i}, \Omega_i, \boxed{P(B|Q)}, p_{T_i}) = \\ P_s(m_i) \cdot P_s(\Omega_i, t_i, \boxed{P(B|Q)}, \sigma_{t_i}) \cdot P_s(\sigma_{t_i}) \\ \cdot P_s(\boxed{P(B|Q)}) \cdot A(\Omega_i, p_{T_i}) \cdot P_s(p_{T_i}). \end{aligned}$$


▶ Muon and Electron Tagging

- ▶ $b \rightarrow l$ transitions are clean tagging method
- ▶ $b \rightarrow c \rightarrow l$ and neutral B-meson oscillations dilute the tagging

▶ Jet-Charge


- ▶ information from tracks in b-tagged jet, when no lepton is found

▶ Calibration using $B^\pm \rightarrow J/\psi K^\pm$



Calibration using $B^\pm \rightarrow J/\psi K^\pm$ events (real data)

- ▶ self tagging non oscillating channel
- ▶ Di-muon candidates in range $2.8 < m(\mu\mu) < 3.4$ GeV
- ▶ $p_T(\mu) > 4$ GeV, $p_T(K^\pm) > 1$ GeV
- ▶ Invariant mass in range $5.0 < m(\mu\mu K^\pm) < 5.6$ GeV
- ▶ $\tau(B) > 0.2$ ps - to reduce prompt component of the combinatorial background



- ▶ Opposite side lepton or jet, with tracks in cone $\Delta R < 0.5$

$$Q = \frac{\sum_i^{N_{\text{tracks}}} q^i(p_T^i)_\kappa}{\sum_i^{N_{\text{tracks}}} q^i(p_T^i)_\kappa} \rightarrow P(Q|B^\pm) \quad Q \in <-1; 1>$$

- The probability to tag a B_s^0 meson as containing a \bar{b} -quark:



$$P(B|Q) = \frac{P(Q|B^+)}{P(Q|B^+) + P(Q|B^-)}$$

Tag method	Efficiency [%]	Effective Dilution [%]	Tagging Power [%]
Tight muon	4.50 ± 0.01	43.8 ± 0.2	0.862 ± 0.009
Electron	1.57 ± 0.01	41.8 ± 0.2	0.274 ± 0.004
Low- p_T muon	3.12 ± 0.01	29.9 ± 0.2	0.278 ± 0.006
Jet	5.54 ± 0.01	20.4 ± 0.1	0.231 ± 0.005
Total	14.74 ± 0.02	33.4 ± 0.1	1.65 ± 0.01

- **Efficiency:** Fraction of signals with specific tagger, $\varepsilon = \frac{N_{\text{tagged}}}{N_{\text{Bcand}}}$
- **Dilution:** $D = (1 - 2w)$, where w is the miss-tag probability
- **Tagging Power:** figure of merit of tagger performance
 - Depends on dilution and efficiency:






$$TP = \varepsilon D^2 = \varepsilon(1 - 2w)^2$$

Tagging performance



Projection and results of the mass-lifetime-angular fit


Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.068	0.038	0.018
$\Delta\Gamma_s$ [ps ⁻¹]	0.067	0.005	0.002
Γ_s [ps ⁻¹]	0.669	0.001	0.001
$ A_{ }(0) ^2$	0.219	0.002	0.002
$ A_0(0) ^2$	0.517	0.001	0.004
$ A_S(0) ^2$	0.046	0.003	0.004
δ_{\perp} [rad]	2.946	0.101	0.097
$\delta_{ }$ [rad]	3.267	0.082	0.201
$\delta_{\perp} - \delta_S$ [rad]	-0.220	0.037	0.010




Combination of the results with the previous from Run 1

- ▶ A Best Linear Unbiased Estimate (BLUE) combination is performed to combine the current result with the Run 1 measurement
- ▶ The BLUE combination uses the measured values and uncertainties of the parameters as well as the correlations between them

Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.076	0.034	0.019
$\Delta\Gamma_s$ [ps ⁻¹]	0.068	0.004	0.003
Γ_s [ps ⁻¹]	0.669	0.001	0.001
$ A_{ }(0) ^2$	0.220	0.002	0.002
$ A_0(0) ^2$	0.517	0.001	0.004
$ A_S ^2$	0.043	0.004	0.004
δ_{\perp} [rad]	3.075	0.096	0.091
$\delta_{ }$ [rad]	3.295	0.079	0.202
$\delta_{\perp} - \delta_S$ [rad]	-0.216	0.037	0.010



- ▶ Analysis of the 2015+2016+2017 ATLAS data performed
- ▶ Results combined with Run1 results
- ▶ Compatible with LHCb and CMS and the SM prediction
- ▶ Complete Run2 analysis ongoing (60 fb⁻¹ more data)




<https://cds.cern.ch/record/2668482>

<https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CONFNOTES/ATLAS-CONF-2019-009/>

Summary and updated overview of the experimental results

- ▶ Analysis of the 2015+2016+2017 ATLAS data performed
- ▶ Results combined with Run1 results
- ▶ Compatible with LHCb and CMS and the SM prediction
- ▶ Complete Run2 analysis ongoing (60 fb⁻¹ more data)




<https://cds.cern.ch/record/2668482>

<https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CONFNOTES/ATLAS-CONF-2019-009/>

Thanks for your attention!

k	$O^{(k)}(t)$	$g^{(k)}(\theta_T, \psi_T, \phi_T)$
1	$\frac{1}{2} A_0(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$2 \cos^2 \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$
2	$\frac{1}{2} A_{\parallel}(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T (1 - \sin^2 \theta_T \sin^2 \phi_T)$
3	$\frac{1}{2} A_{\perp}(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T \sin^2 \theta_T$
4	$\frac{1}{2} A_0(0) A_{\parallel}(0) \cos \delta_{\parallel}$ $\left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin^2 \theta_T \sin 2\phi_T$
5	$ A_{\parallel}(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos(\delta_{\perp} - \delta_{\parallel}) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin(\delta_{\perp} - \delta_{\parallel}) \cos(\Delta m_s t) - \cos(\delta_{\perp} - \delta_{\parallel}) \cos \phi_s \sin(\Delta m_s t)) \right]$	$-\sin^2 \psi_T \sin 2\theta_T \sin \phi_T$
6	$ A_0(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos \delta_{\perp} \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin \delta_{\perp} \cos(\Delta m_s t) - \cos \delta_{\perp} \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin 2\theta_T \cos \phi_T$
7	$\frac{1}{2} A_S(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{2}{3} (1 - \sin^2 \theta_T \cos^2 \phi_T)$
8	$\alpha A_S(0) A_{\parallel}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \sin(\delta_{\parallel} - \delta_S) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos(\delta_{\parallel} - \delta_S) \cos(\Delta m_s t) - \sin(\delta_{\parallel} - \delta_S) \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin^2 \theta_T \sin 2\phi_T$
9	$\frac{1}{2} \alpha A_S(0) A_{\perp}(0) \sin(\delta_{\perp} - \delta_S)$ $\left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin 2\theta_T \cos \phi_T$
10	$\alpha A_0(0) A_S(0) \left[\frac{1}{2}(e^{-\Gamma_H^{(s)} t} - e^{-\Gamma_L^{(s)} t}) \sin \delta_S \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos \delta_S \cos(\Delta m_s t) + \sin \delta_S \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{4}{3} \sqrt{3} \cos \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$



- ▶ Systematics assumed uncorrelated $\rightarrow \text{Total} = \sqrt{\sum_i \text{syst}_i^2}$
- ▶ Tagging systematics dominant for ϕ_s
 - ▶ Accounting for pile-up dependence, calibration curves model and MC precision, "Punzi" PDFs variations, difference between B^\pm and B_s^0 kinematics
- ▶ Fit-model time resolution systematics dominant for Γ_s and $\Delta\Gamma_s$

	ϕ_s [rad]	$\Delta\Gamma_s$ [ps ⁻¹]	Γ_s [ps ⁻¹]	$ A_{ }(0) ^2$	$ A_0(0) ^2$	$ A_S(0) ^2$	δ_\perp [rad]	$\delta_{ }$ [rad]	$\delta_\perp - \delta_S$ [rad]
Tagging	1.7×10^{-2}	0.4×10^{-3}	0.3×10^{-3}	0.2×10^{-3}	0.2×10^{-3}	2.3×10^{-3}	1.9×10^{-2}	2.2×10^{-2}	2.2×10^{-3}
Acceptance	0.7×10^{-3}	$< 10^{-4}$	$< 10^{-4}$	0.8×10^{-3}	0.7×10^{-3}	2.4×10^{-3}	3.3×10^{-2}	1.4×10^{-2}	2.6×10^{-3}
ID alignment	0.7×10^{-3}	0.1×10^{-3}	0.5×10^{-3}	$< 10^{-4}$	$< 10^{-4}$	$< 10^{-4}$	1.0×10^{-2}	7.2×10^{-3}	$< 10^{-4}$
S-wave phase	0.2×10^{-3}	$< 10^{-4}$	$< 10^{-4}$	0.3×10^{-3}	$< 10^{-4}$	0.3×10^{-3}	1.1×10^{-2}	2.1×10^{-2}	8.3×10^{-3}
Background angles model:									
Choice of fit function	1.8×10^{-3}	0.8×10^{-3}	$< 10^{-4}$	1.4×10^{-3}	0.7×10^{-3}	0.2×10^{-3}	8.5×10^{-2}	1.9×10^{-1}	1.8×10^{-3}
Choice of p_T bins	1.3×10^{-3}	0.5×10^{-3}	$< 10^{-4}$	0.4×10^{-3}	0.5×10^{-3}	1.2×10^{-3}	1.5×10^{-3}	7.2×10^{-3}	1.0×10^{-3}
Choice of mass interval	0.4×10^{-3}	0.1×10^{-3}	0.1×10^{-3}	0.3×10^{-3}	0.3×10^{-3}	1.3×10^{-3}	4.4×10^{-3}	7.4×10^{-3}	2.3×10^{-3}
Dedicated backgrounds:									
B_d^0	2.3×10^{-3}	1.1×10^{-3}	$< 10^{-4}$	0.2×10^{-3}	3.1×10^{-3}	1.4×10^{-3}	1.0×10^{-2}	2.3×10^{-2}	2.1×10^{-3}
Λ_b	1.6×10^{-3}	0.4×10^{-3}	0.2×10^{-3}	0.5×10^{-3}	1.2×10^{-3}	1.8×10^{-3}	1.4×10^{-2}	2.9×10^{-2}	0.8×10^{-3}
Fit model:									
Time res. sig frac	1.4×10^{-3}	1.1×10^{-3}	$< 10^{-4}$	0.5×10^{-3}	0.6×10^{-3}	0.6×10^{-3}	1.2×10^{-2}	3.0×10^{-2}	0.4×10^{-3}
Time res. p_T bins	3.3×10^{-3}	1.4×10^{-3}	0.1×10^{-2}	$< 10^{-4}$	$< 10^{-4}$	0.5×10^{-3}	6.2×10^{-3}	5.2×10^{-3}	1.1×10^{-3}
Total	1.8×10^{-2}	0.2×10^{-2}	0.1×10^{-2}	0.2×10^{-2}	0.4×10^{-2}	0.4×10^{-2}	9.7×10^{-2}	2.0×10^{-1}	0.1×10^{-1}