



Uvod v analizo podatkov v eksperimentalni fiziki osnovnih delcev

doc. dr. Marko Bračko

Univerza v Mariboru, FKKT

in



Institut Jožef Stefan




F9 – Odsek za eksperimentalno fiziko osnovnih delcev

IMC2023 – Belle II, IJS, Ljubljana



23. marca 2023



Izmerjeni signali (količine) → vizualizacija




Izmerjeni signali (količine) → vizualizacija



- Za (ročno) pregledovanje je podatkov preprosto preveč ...
- Želimo določiti lastnosti delcev, izmeriti razne vrednosti, konstante, ...
- Potrebujemo izračunane količine, statistične metode, ...

... in kako snov vidimo/razumemo danes?




Osnovni gradniki snovi :

		LEPTONS		QUARKS	
All ordinary particles belong to this group These particles existed just after the Big Bang. Now they are found only in cosmic rays and accelerators	FIRST FAMILY	Electron Responsible for electricity and chemical reactions; it has a charge of -1 e	Electron neutrino Particle with no electric charge, and possibly no mass; billions fly through your body every second ν_e	Up Has an electric charge of plus two-thirds; protons contain two, neutrons contain one u	Down Has an electric charge of minus one-third; protons contain one, neutrons contain two d
	SECOND FAMILY	Muon A heavier relative of the electron; it lives for two-millionths of a second μ	Muon neutrino Created along with muons when some particles decay ν_μ	Charm A heavier relative of the up; found in 1974 c	Strange A heavier relative of the down; found in 1964 s
	THIRD FAMILY	Tau Heavier still; it is extremely unstable. It was discovered in 1975 τ	Tau neutrino not yet discovered but believed to exist ν_τ	Top Heavier still t	Bottom Heavier still; measuring bottom quarks is an important test of electroweak theory b

- vsak delec iz tabele ima svoj anti-delec, npr. e^- in e^+
 - leptoni so nesestavljeni
 - kvarki sestavljajo težje delce - hadrone, npr.
- $p = uud$ ($+2/3e_0 + 2/3e_0 - 1/3e_0$) { e_0 je osnovni naboj: }
 $n = udd$ ($+2/3e_0 - 1/3e_0 - 1/3e_0$) { $\sim 1,6 \cdot 10^{-19} \text{ As}$ }

... in kako snov vidimo/razumemo danes?



gluoni (močna sila)

fotoni (EM sila)

bozoni W^+ , W^- , Z^0 (šibka sila)

gravitoni (gravitacija)

Higgsovo polje (masa delcev) :

Higgsov bozon

Barioni in mezoni: vezana stanja kvarkov in antikvarkov

Barioni

proton: uud

nevtron: udd

Λ : uds

masa

$1 m_p$

$\sim 1 m_p$

$1.2 m_p$

Mezoni

π^+ : kvark u + antikvark \bar{d}

K_S : kvark d + antikvark \bar{s}

J/ψ : kvark c + antikvark \bar{c}

B^0 : kvark d + antikvark \bar{b}

masa

$1/7 m_p$

$1/2 m_p$

$3 m_p$

$5.5 m_p$

In seveda še mnogi drugi delci ...


Kako opazujemo majhne delce?

Kvantna mehanika:
delci se obnašajo kot valovanje
Max Planck

$$E = h\nu = h (c / \lambda)$$


velika energija → velika frekvenca
→ majhna valovna dolžina

Posebna teorija relativnosti:
energija in masa sta izmenljivi
Albert Einstein
 $E=mc^2$




$$E = h\nu$$

$$E = mc^2$$




Sedaj pa še nekaj enačb:

The image shows the equation $E = mc^2$ in white serif font, centered over a background of white, fluffy clouds against a clear blue sky. The sun is visible in the upper right corner, creating a lens flare effect.
$$E = mc^2$$

To je seveda samo poseben primer splošnejše enačbe...


Sedaj pa še nekaj enačb:




$E = mc^2$

To je seveda samo poseben primer splošnejše enačbe...

zato bi morali v resnici zapisati tole:



$E = \gamma mc^2$



$$\gamma = \frac{1}{\sqrt{1 - (v/c)^2}}$$

$\gamma = 1$

$E = mc^2$

ONLY CORRECT
IF AN OBJECT IS
NOT MOVING
WITH RESPECT
TO YOU

Sedaj pa še nekaj enačb: faktor gama (γ)

$$\gamma = \frac{E}{mc^2}$$



$$\gamma = \frac{E \text{ (moving)}}{E \text{ (not moving)}}$$

Ta faktor se torej povečuje, ko pospešujemo delce.

Isti faktor nastopa tudi v definiciji gibalne količine:


$$p = \gamma \times m \times v$$
$$m_{\text{relativistic}} = \gamma m$$
$$p = m_{\text{relativistic}} v$$

S faktorjem γ lahko tudi predefiniramo maso delca.

Prave (relativistične) enačbe :

$$E = \gamma mc^2$$

$$p = \gamma \times m \times v$$



$$E^2 = (pc)^2 + (mc^2)^2$$

E - energy

m - mass

c - speed of light


p - momentum

Z definicijo faktorja γ lahko to enačbo preverite sami...

Prave (relativistične) enačbe :

$$E = \gamma mc^2$$

$$p = \gamma \times m \times v$$



$$E^2 = (pc)^2 + (mc^2)^2$$

E - energy

m - mass

c - speed of light

p - momentum

Dva posebna primera :

Za $p = 0$ (delec, ki miruje) :

$$E = mc^2$$

Za $m = 0$ (foton – γ) :

$$E = pc$$

Invariantna masa – eksperimentalna količina :

$$m c^2 = \sqrt{E^2 - p^2 c^2} \quad (1)$$

Maso delca lahko izrazimo (preverite!) iz prejšnje enačbe... in v idealnem primeru bi bila masa (1) natanko enaka masi delca

Neidealni primer - komplikacije :

- Delci razpadajo → uporabiti moramo razpadne produkte;
- Tudi če zaznamo/izmerimo vse razpadne produkte, povsem natančne meritve niso mogoče → masa (1) je torej \approx enaka masi delca
- Ozadje (npr. podobni razpadi) → še več razlik med maso (1) in maso delca ...

→ Invariantna masa delca :

$$m c^2 = \sqrt{\left(\sum_i E_i\right)^2 - \left(\sum_i \vec{p}_i c\right)^2}$$

$$E_i = \sqrt{\left(\sum_i m_i c^2\right)^2 + \left(\sum_i \vec{p}_i c\right)^2}$$

Indeks i teče po razpadnih produktih delca!

Primer (prave) meritve z detektorjem Belle :

\mathcal{B} Observation of $Z^+(4430)$ state

PRL 100, 142001(2008)
657 $B\bar{B}$

$Z(4430)^+ \rightarrow \psi(2S)\pi^+$:


Charged state that decays like charmonium (= charged charmonium-like state)

$$\text{Br}(\bar{B}^0 \rightarrow K^- Z^+(4430)) \times \text{Br}(Z^+(4430) \rightarrow \pi^+ \psi') = (4.1 \pm 1.0 \pm 1.4) \times 10^{-5}$$

Necessary ingredients:

- Histogram \rightarrow bins of $m_{\text{invariant}}$
- Just specific decay modes are used
- Errors from measurements ...
- Background:
 - wrong part. identification
 - missing particles in reconstr.
 - wrong combinations/decay modes
 - ...

\rightarrow First serious tetraquark candidate



Primer (prave) meritve z detektorjem Belle :


\mathcal{B} Observation of $Z^+(4430)$ state

PRL 100, 142001(2008)
657 $B\bar{B}$

$Z(4430)^+ \rightarrow \psi(2S)\pi^+$:

Charged state that decays like charmonium (= charged charmonium-like state)

$$\text{Br}(\bar{B}^0 \rightarrow K^- Z^+(4430)) \times \text{Br}(Z^+(4430) \rightarrow \pi^+ \psi') = (4.1 \pm 1.0 \pm 1.4) \times 10^{-5}$$



➔ First serious tetraquark candidate

Primer (prave) meritve z detektorjem Belle :

Observation of $Z^+(4430)$ state

PRL 100, 142001(2008)
657 $B\bar{B}$


$Z(4430)^+ \rightarrow \psi(2S)\pi^+$:

Charged state that decays like charmonium (= charged charmonium-like state)

$$\text{Br}(\bar{B}^0 \rightarrow K^- Z^+(4430)) \times \text{Br}(Z^+(4430) \rightarrow \pi^+ \psi') = (4.1 \pm 1.0 \pm 1.4) \times 10^{-5}$$

Necessary ingredients:

- Histogram \rightarrow bins of m_{invar}
- Just specific decay modes are used (statistics)
- Errors from measurements ...
- Background:
 - wrong part. identification
 - missing particles in reconstr.
 - wrong combinations/decay modes
 - ...



- Extract the parameters: fitting the data distribution with functions

Primer komplikacij : različni razpadni načini delcev

Properties of kaons

Particle name	Particle symbol	Antiparticle symbol	Quark content	Rest mass (MeV/c ²)	I ^G	J ^{PC}	S	C	B'	Mean lifetime (s)	Commonly decays to (>5% of decays)
Kaon ^[1]	K ⁺	K ⁻	u \bar{s}	493.677 ± 0.016	1/2	0 ⁻	1	0	0	(1.2380 ± 0.0021) × 10 ⁻⁸	$\mu^+ + \nu_\mu$ or $\pi^+ + \pi^0$ or $\pi^+ + \pi^+ + \pi^-$ or $\pi^0 + e^+ + \nu_e$
Kaon ^[2]	K ⁰	\bar{K}^0	d \bar{s}	497.611 ± 0.013	1/2	0 ⁻	1	0	0	[a]	[a]
K-Short ^[3]	K _S ⁰	Self	$\frac{d\bar{s} + s\bar{d}}{\sqrt{2}}$ [b]	497.611 ± 0.013 ^[c]	1/2	0 ⁻	(*)	0	0	(8.954 ± 0.004) × 10 ⁻¹¹	$\pi^+ + \pi^-$ or $\pi^0 + \pi^0$
K-Long ^[4]	K _L ⁰	Self	$\frac{d\bar{s} - s\bar{d}}{\sqrt{2}}$ [b]	497.611 ± 0.013 ^[c]	1/2	0 ⁻	(*)	0	0	(5.116 ± 0.021) × 10 ⁻⁸	$\pi^+ + e^- + \nu_e$ or $\pi^+ + \mu^- + \nu_\mu$ or $\pi^0 + \pi^0 + \pi^0$ or $\pi^+ + \pi^0 + \pi^-$



$$D^{*+} \rightarrow D^0 \pi^+$$

$$D^{*-} \rightarrow D^0 \pi^-$$


$$D^0 \rightarrow K^+ \pi^- / K^- \pi^+$$

Možne izboljšave :

Analysis tools: B-meson selection



Reconstructing B meson decays at Y(4S):
use two variables,
beam-constrained mass M_{bc}
(energy-substituted mass m_{ES})
and
energy difference ΔE



$$\Delta E \equiv \sum E_i^{CMS} - E_{beam}^{CMS}$$

$$M_{bc} = \sqrt{(E_{beam}^{CMS})^2 - (\sum \vec{p}_i^{CMS})^2}$$

Zaključek :

Sedaj ste (vsaj v grobem) pripravljeni, da tudi sami poskusite priti do rezultatov ...

Srečno!

